

KPL Cy™3 - Labeled Antibody to Rabbit IgG (H+L)

Produced in Goat

Catalog No.

5230-0359 (072-01-15-06)

Size

1.0 mg

DESCRIPTION

Affinity purified antibody isolated from a pool of serum from goats immunized with purified rabbit IgG was labeled with Cy3 using a proprietary method. Cy conjugates provide increased brightness and greater photostability than other fluorophores (ex. – FITC). This conjugate is designed for use in immunofluorescence assays, including flow cytometry, immunohistology, Western blotting and microarrays.

FORM/STORAGE

Lyophilized. Store at 2-8°C until rehydrated. Stable for a minimum of 1 year when stored at 2-8°C.

STABILIZER AND PRESERVATIVE

IgG free bovine serum albumin (BSA) is added as a protein stabilizer. Non-sterile.

ANTIBODY CONCENTRATION

The quantity of affinity purified antibody is 1.0 mg as determined by UV absorbance at 280 nm. Upon rehydration with water, the solution will contain 1% BSA, 100 mM phosphate, 150 mM sodium chloride, 0.02% sodium azide, pH 7.4. The conjugate will be at a concentration of 1.0 mg/mL.

F/P RATIO

Molar fluorophore/antibody protein ratio is from 4:1 – 7:1.

EXCITATION/EMISSION VALUES

Cy3 is excited at 550 nm and emits at 570 nm.

SPECIFICITY/CROSS REACTIVITY

Tested by gel diffusion and ELISA techniques as applicable. This product reacts specifically with rabbit IgG and may recognize other immunoglobulin types that have light chains in common with IgG. Reactivity to IgG subclasses has not been tested. Antibodies to rabbit IgG may cross-react with immunoglobulins of other mammalian species if common binding sites are shared.

REHYDRATION AND STORAGE

Note: Rehydration of antibodies in buffers other than those listed here is not recommended.

Rehydration: Rehydrate with 1 mL of reagent quality water. Rotate the vial until the lyophilized pellet is totally dissolved.

Use: Prior to use, dilute to desired concentration with PBS or 1% BSA Diluent/Blocking Solution. The diluted conjugate should be used immediately.

Storage: The rehydrated product is stable for a minimum of 1 year at 2-8°C.

SUGGESTED WORKING DILUTIONS

Optimal working concentrations should be determined experimentally for each application. Prepare working dilution in PBS or other buffer such as KPL BSA Diluent/Blocking Solution. Dilution is not recommended for long-term storage. Suggested starting dilutions are listed below:

Histo/Cytochemical/Flow Cytometry:

1:100 – 1:1,000 10 µg/mL – 1 µg/mL

PRODUCT SAFETY AND HANDLING

This product is considered non-hazardous as defined by The Hazard Communication Standard (29 CFR 1910.1200). Avoid contact with skin and eyes. In case of contact or spillage, clean with copious amounts of water. Dispose of via institutional guidelines.

TRADEMARKS

Cy™ is a trademark of GE Healthcare, Inc.

RELATED PRODUCTS

	CAT. NO.
KPL BSA Diluent/Blocking Solution	5140-0006 (50-61-00)
KPL Coating Solution	5150-0014 (50-84-00)
KPL Wash Solution	5150-0008 (50-63-00)
KPL Fluorescent Mounting Medium	5540-0002 (71-00-06)
KPL 10X PBS	5460-0022 (51-13-01)

The product listed herein is for research use only and is not intended for use in human or clinical diagnosis.